

Faraci Lange - Vaccine Breakdown

| Vaccine | Illness, Disability, or Injury Covered | Time Period for First Symptom After Vaccine Administration |
|---|--|--|
| Vaccines containing tetanus toxoid (e.g., DTaP, DTP, DT, Td, or TT) | Anaphylaxis or anaphylactic shock | 4 hours |
| | Brachial Neuritis | 2-28 days |
| | Shoulder Injury Related to Vaccine Administration | ≤48 hours |
| Vaccines containing whole cell pertussis bacteria, extracted or partial cell pertussis bacteria, or specific pertussis antigen(s) (e.g., DTP, DTaP, P, DTP-Hib) | Anaphylaxis or anaphylactic shock | 4 hours |
| | Encephalopathy (or encephalitis) | 72 hours |
| | Shoulder Injury Related to Vaccine Administration | ≤48 hours |
| Measles, mumps, and rubella vaccine or any of its components (e.g., MMR, MR, M, R) | Anaphylaxis or anaphylactic shock | 4 hours |
| | Encephalopathy (or encephalitis) | 5-15 days |
| | Shoulder Injury Related to Vaccine Administration | ≤48 hours |
| Vaccines containing rubella virus (e.g., MMR, MR, R) | Chronic arthritis | 7-42 days |
| Vaccines containing measles virus (e.g., MMR, MR, R) | Thrombocytopenic purpura | 7-30 days |
| | Vaccine-Strain Measles Viral Infection in an immunodeficient recipient | 6 months |
| Vaccines containing polio live virus (OPV) | Paralytic Polio | 30 days-6 months |
| | Vaccine-Strain Polio Viral Infection | 30 days-6 months |
| Vaccines containing polio inactivated virus (e.g., IPV) | Anaphylaxis or anaphylactic shock | 4 hours |
| | Shoulder Injury Related to Vaccine Administration | ≤48 hours |
| Hepatitis B. vaccines | Anaphylaxis or anaphylactic shock | 4 hours |
| | Shoulder Injury Related to Vaccine Administration | ≤48 hours |
| Seasonal influenza vaccines | Anaphylaxis or anaphylactic shock | ≤4 hours |
| | Shoulder Injury Related to Vaccine Administration | ≤48 hours |
| | Guillain-Barré Syndrome | 3-42 days |